Oswestry Disability Index

Pain prevents me from standing at all.

Sed	ction 1 – Pain Intensity		My sleep is never disturbed by pain.
			My sleep is occasionally disturbed by pain.
	I have no pain at the moment.		Because of pain, I have less than 6 hours sleep.
	The pain is very mild at the moment.		Because of pain, I have less than 4 hours sleep.
	The pain is moderate at the moment.		Because of pain, I have less than 2 hours sleep.
	The pain is fairly severe at the moment.		Pain prevents me from sleeping at all.
	The pain is very severe at the moment.		· ·
	The pain is the worst imaginable at the moment.	Sec	ction 8 – Sex life (if applicable)
Sec	ction 2 – Personal Care (washing, dressing, etc.)		My sex life is normal and causes no extra pain.
			My sex life is normal but causes some extra pain.
	I can look after myself normally but it is very painful.		My sex life is nearly normal but is very painful.
	I can look after myself normally but it is very painful.		My sex life is severely restricted by pain.
	It is painful to look after myself and I am slow and careful.		My sex life is nearly absent because of pain.
	I need some help but manage most of my personal care.		Pain prevents any sex life at all.
	I need help every day in most aspects of my personal care.		
	I need help every day in most aspects of self-care.	Sec	ction 9 – Social Life
	I do not get dressed, wash with difficulty, and stay in bed.		
_	Too not got allocood, mach min amounty, and only in soul		My social life is normal and cause me no extra pain.
Sed	ction 3 - Lifting	_	My social life is normal but increases the degree of pain.
	· ·	_	Pain has no significant effect on my social life apart from limitingmy
	I can lift heavy weights without extra pain.	_	more energetic interests, i.e. sports.
	I can lift heavy weights but it gives extra pain.		- · · · · · · · · · · · · · · · · · · ·
_	Pain prevents me from lifting heavy weights off the floor, but I can		Pain has restricted my social life and I do not go out as often.
_	manage if they are conveniently positioned (i.e. on a table).		Pain has restricted social life to my home.
	Pain prevents me from lifting heavy weights, but I can manage light to		I have no social life because of pain.
_		Sec	ction 10 – Traveling
	medium weights if they are conveniently positioned.		g
	I can lift only very light weights.		I can travel anywhere without pain.
_	I cannot lift or carry anything at all.	_	I can travel anywhere but it gives extra pain.
Sec	ction 4 – Walking	_	Pain is bad but I manage journeys of over two hours.
-	numing		Pain restricts me to short necessary journeys under 30 minutes.
	Pain does not prevent me walking any distance.	_	Pain prevents me from traveling except to receive treatment.
	Pain prevents me walking more than 1mile.	_	and prevents the norm traveling except to receive treatment.
_	Pain prevents me walking more than ¼ of a mile.	Soc	ction 11 - Previous Treatment
_	Pain prevents me walking more than 100 yards.	360	Citori II - Frevious Treatment
_	I can only walk using a stick or crutches.	0	
_	I am in bed most of the time and have to crawl to the toilet.		r the past three months have you received treatment, tablets or
_	Tain in sec most of the time and have to craw to the tollet.		dicines of any kind for your back or leg pain? Please check the ropriate box.
Sed	ction 5 – Sitting		No
		_	Yes (if yes, please state the type of treatment you have received)
	I can sit in any chair as long as I like.	_	Too (ii you, ploade state the type of troublett you have received)
	I can sit in my favorite chair as long as I like.		
	Pain prevents me from sitting for more than 1 hour.		
	Pain prevents me from sitting for more than ½ hour.		
	Pain prevents me from sitting for more than 10		
	minutes.		
	Pain prevents me from sitting at all.		
Sec	ction 6 – Standing		
	I can stand as long as I want without extra pain.		
	I can stand as long as I want but it gives me extra pain.		
	Pain prevents me from standing more than 1 hour.		
	Pain prevents me from standing for more than ½ an hour.		
	Pain prevents me from standing for more than 10 minutes		

Section 7 - Sleeping

Neck Disability Index

I cannot concentrate at all. (5)

This questionnaire has been designed to give the doctor information as to how your neck pain has affected your ability to manage in everyday life. Please answer every section and mark in each section only the ONE box which applies to you. We realize you may consider that two of the statements in any one section relate to you, but please just mark the box which most closely describes your problem.

Section 7 - Work

Sec	ction 1 – Pain Intensity		I can do as much work as I want to. (0)
	I have no pain at the moment. (0)		I can do my usual work, but no more. (1)
	The pain is very mild at the moment. (1)		I can do most of my usual work, but no more. (2)
	The pain is moderate at the moment. (2)		I cannot do my usual work. (3)
	The pain is fairly severe at the moment. (3)		I can hardly do any work at all. (4)
	The pain is very severe at the moment. (4)		I cannot do any work at all. (5)
	The pain is the worst imaginable at the moment. (5)		
		Sect	ion 8 – Driving
Sec	ction 2 - Personal Care (Washing, Dressing, etc.)		I can drive my car without any neck pain. (0)
	I can look after myself normally without causing extra pain. (0)		I can drive my car as long as I want with slight pain in my neck. (1)
	I can look after myself normally but it causes extra pain. (1)		I can drive my car as long as I want with moderate pain in my neck. (2
	It is painful to look after myself and I am slow and careful. (2)		I cannot drive my car as long as I want because of moderate pain in
	I need some help but manage most of my personal care. (3)		my neck. (3)
	I need help every day in most aspects of self care. (4)		I can hardly drive at all because of severe pain in my neck. (4)
	I do not get dressed, I wash with difficulty and stay in bed. (5)		I cannot drive my car at all. (5)
Sec	etion 3 – Lifting	Sect	ion 9 – Sleeping
	I can lift heavy weights without extra pain. (0)		I have no trouble sleeping. (0)
	I can lift heavy weights but it gives extra pain. (1)		My sleep is slightly disturbed (less than 1 hour sleepless). (1)
	Pain prevents me from lifting heavy weights off the floor, but I can		My sleep is mildly disturbed (1-2 hours sleepless). (2)
	manage if they are conveniently positioned, for example on a table. (2)		My sleep is moderately disturbed (2-3 hours sleepless). (3)
	Pain prevents me from lifting heavy weights, but I can manage light to		My sleep is greatly disturbed (3-5 hours sleepless). (4)
	medium weights if they are conveniently positioned. (3)		My sleep is completely disturbed (5-7 hours sleepless). (5)
	I can lift very light weights. (4)		
	I cannot lift or carry anything at all. (5)	Sect	ion 10 – Recreation
			I am able to engage in all my recreation activities with no neck pain at
Sec	ction 4 - Reading		all. (0)
	I can read as much as I want to with no pain in my neck. (0)		I am able to engage in all my recreation activities, with some pain in
	I can read as much as I want to with slight pain in my neck. (1)		my neck. (1)
	I can read as much as I want with moderate pain in my neck. (2)		I am able to engage in most, but not all, of my usual recreation
	I cannot read as much as I want because of moderate pain in my neck.		activities because of pain in my neck. (2)
	(3)		I am able to engage in a few of my usual recreation activities because
	I can hardly read at all because of severe pain in my neck. (4)		of pain in my neck. (3)
	I cannot read at all. (5)		I can hardly do any recreation activities because of pain in my neck.
			(4)
Sec	ction 5 – Headaches		I cannot do any recreation activities at all. (5)
	I have no headaches at all. (0)		
	I have slight headaches that come infrequently. (1)		
	I have moderate headaches which come infrequently. (2)		
	I have moderate headaches which come frequently. (3)		
	I have severe headaches which come frequently. (4)	0-4	No disability
	I have headaches almost all the time. (5)	5-14	Mild disability
		15-2	4 Moderate disability
Sec	ction 6 - Concentration	25-3	4 Severe disability
	I can concentrate fully when I want to with no difficulty. (0)	> 35	Complete disability
	I can concentrate fully when I want to with slight difficulty. (1)		
	I have a fair degree of difficulty in concentrating when I want to. (2)		
	I have a lot of difficulty in concentrating when I want to. (3)		
	Lhave a great deal of difficulty in concentrating when I want to (4)		

Functional Strength of the Cervical Spine

Starting Position	Action	Functional Test
Supine lying	Lift head keeping chin tucked in (neck flexion)	6 to 8 repetitions: functional 3 to 5 repetitions: functionally fair 1 to 2 repetitions: functionally poor 0 repetitions: nonfunctional
Prone lying	Lift head backward (neck extensions)	Hold 20 to 25 seconds: functional Hold 10 to 19 seconds: functionally fair Hold 1 to 9 seconds: functionally poor Hold 0 seconds: nonfunctional
Side lying (pillows under head so head is not side flexed)	Life head sideways away from pillow (neck side flexion) (must be repeated or other side)	Hold 20 to 25 seconds: functional Hold 10 to 19 seconds: functionally fair Hold 1 to 9 seconds: functionally poor Hold 0 seconds: nonfunctional
Supine lying	Lift head off bed and rotate to one side keeping head off bed or pillow (neck rotation) (must be repeated both ways)	Hold 20 to 25 seconds: functional Hold 10 to 19 seconds: functionally fair Hold 1 to 9 seconds: functionally poor Hold 0 seconds: nonfunctional

SCORING TECHNIQUE FOR THE OSWESTRY LOW BACK DISABILITY QUESTIONNAIRE AND NECK DISABILITY INDEX

1. Each of the 10 sections is scored separately (0 to 5 points each) and then added up (max. total = 50).

Example:

Section 1. Pain Intensity	Point Value
A I have no pain at the moment	0
B The pain is very mild at the moment	1
C The pain is moderate at the moment	2
D The pain is fairly severe at the moment	3
E The pain is very severe at the moment	4
F The pain is the worst imaginable	5

- 2. If all 10 sections are completed, simply double the patient's score.
- 3. If a section is omitted, divide the patient's total score by the number of sections completed times 5.

Formula:	Patient's Score		
		X 100 =	% DISABILITY
	No. of sections completed x 5	X 100 =	/0 BIG/(BILIT I

Example:

If 9 of 10 sections are completed, divide the patient's score by $9 \times 5 = 45$.

Patient's Score 2

Number of sections completed: $9 (9 \times 5 = 45)$

 $22/45 \times 100 = 48\%$ disability

4. Interpretation of disability scores (from original article):

SCORE INTERPRETATION OF THE OSWESTRY LBP DISABILITY QUESTIONNAIRE		
0-20% Minimal disability	Can cope with most ADLs. Usually no treatment is needed, apart from advice on lifting, sitting, posture, physical fitness, and diet. In this group, some patients have particular difficulty with sitting and this may be important if their occupation is sedentary (typist, driver, etc.)	
20-40% Moderate disability	This group experiences more pain and problems with sitting, lifting, and standing. Travel and social life are more difficult and they may well be off work. Personal care, sexual activity, and sleeping are not grossly affected, and the back condition can usually be managed by conservative means.	
40-60% Severe disability	Pain remains the main problem in this group of patients, but travel, personal care, social life, sexual activity, and sleep are also affected. These patients require detailed investigation.	
Back pain impinges on all aspects of these patients' lives both at home and at work. Positive intervention is required.		
80-100%	These patients are either bed-bound or exaggerating their symptoms. This can be evaluated by careful observation of the patient during the medical examination.	
Data compiled from Fairbanks et al, 1980.		